

## WOUND PRODUCT SELECTION GUIDE

Gippsland		WOUND EXUDATE LEVEL			
DRESSING / TRADE NAME * Primary dressing ** Secondary dressing	DRESSING FUNCTION AND COMMENTS	None	LOW <u>&lt;</u> 7 <sup>days</sup>	MODERATE 2-3 days	HIGH < 24 <sup>hrs</sup>
	Onen mesh estten much viscore		wear	wear	wear
* Non-adherent dressings: Ugotul, Adaptic, Atrauman, Jelonet	Open mesh, cotton, rayon, viscose or gauze impregnated with gel, silicone to form a pliable non-stick interface.		✓	✓	✓
*Low absorbent dressings: Melolin, Primapore, Interpose, Telfa, Post op island dressings	Post op dressings or wounds almost healed. For protection in healed wounds.	$\checkmark$	$\checkmark$		
<b>**Semi permeable films:</b> Opsite, Tegaderm, Mepitel film	Reduces wound pain Caution on removal to avoid skin tear (trauma)	~	✓		
* <b>Hydrogels:</b> Instrasite, Solosite, Duoderm gel, Purilon gel	Autolytic debridement. Pain reduction. Requires a secondary dressing	✓	$\checkmark$		
* <b>Hydrofibres:</b> Aquacel, Durafibre, Activheal, Aquafibre	Ideal for packing wounds Requires moisture to be effective.			$\checkmark$	$\checkmark$
*Alginates: Kaltostat, Algisite M, Sorbsan, Melgisorb	Do not use on dry wounds. Requires moisture to be effective. <i>Nb. Kaltostat is a haemostat.</i>		✓	$\checkmark$	
<b>**Foams:</b> <i>Mepilex, Allevyn,</i> <i>Biatain, Activheal</i>	With or without borders Some use silicone Can be used as a primary or secondary dressing		✓	$\checkmark$	$\checkmark$
<b>**Hydrocolloids:</b> Comfeel, Duoderm, Replicare	Maintains moisture on healing wounds. Can be used as a primary or secondary dressing	✓	$\checkmark$		
<b>**Superabsorbent polymers:</b> <i>Exudry, Mextra, Relevo, Zetuvit</i>	Inner dressing contains superabsorbent particles Secondary dressing			$\checkmark$	$\checkmark$
*Silvers: Acticoat (3,7 Flex and Moisture control), Aquacel Ag, Melgisorb Ag, Biatain Ag, Mepilex Ag,	Most foams and alginates have a version with a silver component added. Nb. Acticoat and Acticoat 7 need premoistening with sterile water. Blue side to wound		✓	✓	✓
*lodine based: lodosorb (Cadexomer lodine - Ointment, Powder and Flex) Inadine is 10% Povidone lodine	Antimicrobial effect Needs replacing when the colour leaves dressing Not absorbent, needs secondary dressing		✓	✓	✓
* <b>Hydrophobic dressings:</b> Sorbact – Gel sheets, ads, Gauze	Hydrophobic dressing attract bacteria and fungi to dressing Requires secondary dressing		$\checkmark$	$\checkmark$	$\checkmark$
*Hypertonic saline: Mesalt, Curasalt	Causes an osmotic draw to clean wound, necrotic tissue or purulent debris. Requires secondary dressing		$\checkmark$	$\checkmark$	
*Honey: Medihoney, Apinate dressing	Autolytic debridement. Antimicrobial activity Requires secondary dressing		$\checkmark$	$\checkmark$	
*Alginate gels: Flaminal Forte, Flaminal Hydro	Autolytic debridement. Antimicrobial activity Requires secondary dressing		✓		